

Anticoagulants in Older Adults


Anita Rajasekhar MD, MS, FACP

November 3, 2024

Learning Objectives

- Review evidence to support anticoagulation in older adults
- Explain the unique challenges of anticoagulation including increased risks of bleeding, frailty, and comorbid conditions in older adults
- Discuss how to tailor anticoagulation therapy in older adults by applying risk assessment tools to balance bleeding and thrombotic risks
- Evaluate patient cases to differentiate between high-risk and low-risk older adults for anticoagulation, and analyze when to adjust or discontinue therapy based on clinical factors


Roadmap for Our Discussion



Age-related physiologic changes

Age-related physiologic changes	Potential impact on OAC and outcomes
Decreased skeletal muscle mass & total body water	<ul style="list-style-type: none">• Increased plasma concentrations of apixaban and edoxaban if <60kg• Increased risk of major bleeding with edoxaban if body weight \leq60kg• Decreased hepatic clearance of warfarin
Decline in GFR	<ul style="list-style-type: none">• Increased plasma concentrations of dabigatran>edoxaban>rivaroxaban>apixaban, esp if CrCl <30ml/min
Decrease in liver size and blood flow	<ul style="list-style-type: none">• Increased DOAC plasma concentration if moderate (rivaroxaban) or severe (apixaban, edoxaban, dabigatran) hepatic dysfunction• Reduced warfarin clearance
Reduced activity in Vit K redox recycling symptom	<ul style="list-style-type: none">• Increased warfarin sensitivity with about 20% lower warfarin dose requirements
Increased prevalence amyloid angiopathy, and cerebral atrophy	<ul style="list-style-type: none">• Increased risk ICH
Increased prevalence of diverticular and peptic ulcer disease	<ul style="list-style-type: none">• Increased risk of GI bleeding

Geriatric syndromes



Case 1: Audience Response Question



87M to ER after fall



BP 154/85 HR ~95 bpm (irreg), wt 82kg



HTN, DM2, HLD, OA, macular degeneration



Hb 12.0 g/dL, INR 1.0
SCr 1.1mg/dL

- A. Low dose aspirin
- B. LMWH → warfarin
- C. Apixaban
- D. Rivaroxaban
- E. DOAC + ASA





Lisinopril, Metformin,
Atorvastatin, Meloxicam,
low dose aspirin




ECG - atrial fibrillation

What would you start this patient on to reduce the risk of AFib-related stroke?

Why is this important to discuss?



US population is aging



AFib prevalence increases with age



<https://www.census.gov/en.html>
Go et al. JAMA. 2001 May 9;285(18):2370-5.

OCSP: Oxfordshire Community Stroke Project; OXVASC: Oxford Vascular Study. Yiin GS, et al. *Circulation* 2014; 130(15):1236-44.

Stroke causes death and disability



30-day mortality up to 20%



Kelly-Hayes et al, J Stroke Cerebrovasc Dis. 2003 May;12(3):119-26. Grysiwicz Neurol Clin. 2008 Nov;26(4):871-95. Feigin et al, Lancet Neurol. 2003;2(1):43.

Stroke risk assessment in Afib: CHA₂DS₂-VASc score



CHADS ₂ – VASc Score		
C	Congestive Heart Failure	1
H	Hypertension (>140/90 mmHg)	1
A	Age ≥ 75	2
D	Diabetes Mellitus	1
S₂	Prior TIA or stroke	2
V	Vascular disease (MI, aortic plaque etc)	1
A	Age 65-74	1
Sc	Sex category (Female = 1 pt)	1

Score	Risk of stroke
0	0.2% Low
1	0.6% Moderate
2	2.2% High
3	3.2%
4	4.8%
5	7.2%
6	9.7%
7	11.2%
8	10.8%
9	12.2%

No AC

Consider AC

Recommend AC




VTE recurrence: risk assessment models

	Men and HERDOO2	Vienna Risk Model	DASH
Gender	X	X	X
D-dimer	X	X	X
Signs of Post-thrombotic syndrome	X		
Obesity	X		
Age	X		X
Location of DVT/PE		X	
Provoked?			X

RCT Evidence for Anticoagulation in Older Patients with AFib

Network metanalysis of 28,135 patients >75 years with AFib



Stroke/Systemic embolism in age ≥ 75 yo: DOACs vs VKA in ARISTOPHANES registry

	apixaban vs. warfarin		dabigatran vs. warfarin		rivaroxaban vs. warfarin	
	incidence rate/ 100 person-years	HR (95% CI)	incidence rate/ 100 person-years	HR (95% CI)	incidence rate/ 100 person-years	HR (95% CI)
Stroke/SE						
75-79 years	1.03 v. 1.79	0.53 (0.42-0.66)	1.51 v. 1.75	0.86 (0.63-1.17)	1.33 v. 1.72	0.76 (0.64-0.9)
≥ 80 years	1.76 v. 2.59	0.62 (0.55-0.71)	2.14 v. 2.59	0.82 (0.66-1.03)	2.16 v. 2.57	0.79 (0.71-0.88)

Bleeding Risk Also Increases with Age


HAS-BLED Bleed Score	
Clinical characteristic	Points
Hypertension	1
Renal or hepatic dysfunction	1 or 2
History of stroke	1
History of bleeding	1
Labile INR	1
Age >65	1
Drugs or alcohol	1 or 2

DOAC Bleed Score	
Clinical characteristic	Points
Age 65-69	2
Age 70-74	3
Age 75-79	4
Age ≥ 80	5
CrCl 30-60 ml/min	1
CrCl <30 ml/min	2
BMI <18.5 kg/m ²	1
Stroke/TIA/embolism	1
Diabetes	1
Hypertension	1
Single/Dual antiplatelet	2/3
NSAID use	1
Bleeding history	3
Liver disease	2

- Bleed risk scores should NOT be used in isolation to decide on prescribing anticoagulants
- Assess for & address modifiable bleed risk factors
- In our case patient
 - Need for aspirin?
 - Optimize BP
 - Minimize NSAID use
 - Assess fall risk with mac degen



Real world evidence of Major bleeding on DOACs (ARISTOPHANES registry N=466,991)



Major Bleeding in age \geq 75yo: DOACs vs VKA in ARISTOPHANES registry

	apixaban vs. warfarin		dabigatran vs. warfarin		rivaroxaban vs. warfarin	
	incidence rate/ 100 person-years	HR (95% CI)	incidence rate/ 100 person-years	HR (95% CI)	incidence rate/ 100 person-years	HR (95% CI)
Major bleeding						
75-79 years	3.59 v. 5.45	0.61 (0.54-0.69)	3.63 v. 5.34	0.68 (0.56-0.82)	5.94 v. 5.44	1.07 (0.98-1.16)
\geq 80 years	4.84 v. 6.88	0.65 (0.6-0.7)	5.9 v. 6.61	0.89 (0.78-1.02)	8.16 v. 6.52	1.17 (1.1-1.25)

Comment on: 2023 updated AGS Beers Criteria for potentially inappropriate medication use in older adults

Warfarin- avoid as initial therapy (unchanged)

Dabigatran- use with caution (unchanged)

Rivaroxaban- **CHANGED** from use with caution to avoid for long-term treatment

Key excerpts from Beers Criteria

A recommendation to “avoid” is not an absolute contraindication.

Medications listed in the Beer's Criteria are *potentially* inappropriate, not definitely inappropriate.

The criteria are a blunt instrument, and it is not possible to delineate all specialized use cases and possible exceptions to the criteria.

Prescribing for older adults is often a complex endeavor involving consideration of many factors , particularly the preferences and goals of the older person and their family.


Rivaroxaban may be a reasonable option in select circumstances.




Anticoagulation + Antiplatelet Therapy: Common Practice

- 1/3 of patients on OAC are also on inappropriate antiplatelet therapy

Percent of Patients Without Recent Myocardial Infarction or Valve Replacement on Aspirin and Warfarin by Year




Percent of Patients Without Recent Myocardial Infarction or Valve Replacement on Aspirin and DOAC by Year




Anticoagulation + Antiplatelet Therapy: Risk vs Reward

- Addition of antiplatelet potentiates efficacy of OAC?
- No difference in thrombotic outcomes (exception mechanical heart valve)
- 2x increase in major bleeding

Patients with chronic Afib, CAD, high risk CAD, mechanical valve



Patient with Acute VTE




Antiplatelet + Anticoagulant Use in NH residents

- Cross-sectional study
- 12 NH chains (709 facilities across 40 states)
- ≥ 100 days in a NH and had AF and a CHA2DS2-VASc (≥ 1 men, ≥ 2 women)
- Stratified:
 - 1) OAC plus antiplatelets (N=582)
 - 2) OAC only (N=1281)
 - 3) antiplatelets only (N=1523)
 - 4) no antithrombotic (N=1366)

12% receiving dual antithrombotic therapy and 45% receiving antiplatelets with no indication for use

Key Considerations in Choosing an Anticoagulant WITH Your Older Patient



Case 2: Audience Response Question

84-year-old man at his PCP office is diagnosed with new onset AFib.

He recently moved into an assisted-living facility after wife died 6 months ago.

HTN managed x 20 years with ACEI. **Severe OA** causing mobility limitations. **DM2** occasionally requiring medication adjustment. **Early-stage dementia** with mild memory impairment but still able to make decisions about his care.

Endorses mild fatigue and **has fallen once in the last year** (able to get up on his own and did not sustain serious injury)

Weight 72kg (5 kg↓)

SCr 1.2 mg/dL

INR 1.0

CHA₂DS₂-Vasc= 4

What would you start to prevent AFib-related stroke?

- A. Low-dose ASA
- B. LMWH → VKA
- C. DOAC
- D. Withhold antiplatelets and anticoagulation




Older patients less likely to be prescribed OAC

- Swedish registry (2009-2012) of 12,000 first-time stroke patients with AFib

Age group	Valid Observations	OAC Prescribing Frequency	Proportion (%)
18-69	1789	1098	61.4
70-79	2909	1531	52.6
80-89	5342	1551	29.0
90+	1993	209	10.5

Prescriber criteria for withholding AC




Other reasons:

- Short life-expectancy
- Fear of bleeding
- Perceived harm > benefit
- Poor health
- Geriatric syndromes

Net clinical benefit of OACs is higher in older age groups


Any OAC vs. No AC: PREFER-AF (European Registry) 2012-2014





Most OAC treatment in the registry was warfarin (~6% NOACs)

Net clinical benefit of DOACs preserved ≥ 75 years

STROKE RISK*



RISK OF MAJOR BLEEDING



Case 3: Audience Response Question

82-year-old woman on **warfarin x 10 years** for AFib brought into anticoagulation clinic by her grandson

Increasingly reliant on others for help with ADLs, including transportation to medical appointments, because of vision, stamina and cognition issues

Weight 57 kg (5 kg↓) SCr 1.8 mg/dL INR 2.2 (TTR ~75%)

Her grandson asks you about “another blood thinner” he saw on TV that is “easier to use” and asks if this might be an option for her

How would you adjust this patient's anticoagulation regimen?

- A. Switch to low-dose ASA
- B. Switch to rivaroxaban
- C. Switch to apixaban
- D. Continue current therapy with warfarin




Frailty

Increased vulnerability due to aging-associated decline in reserve & function across multiple physiologic systems


Not well represented in RCTs

Multiple frailty scoring tools

Clinicians implicitly weigh multiple risk factors when deciding AC vs. no AC



Meta-analysis of frailty in AFib (N=1,187,000)



Prevalence of Frailty: 40%

Prevalence of Pre-frailty: 35%



OAC under-prescription in frail patients vs prefrail/robust:
OR 0.83 (0.61-1.06)

Frail >80yo → OAC significantly less prescribed

Frail patients at higher risk for major outcomes compared to robust patients



All-cause death
OR 5.56 (3.46-8.94)



Stroke
OR 1.59 (1.00-2.52)



Bleeding
OR 1.64 (1.11-2.41)

FRAIL-AF: Design

Circulation

Volume 149, Issue 4, 23 January 2024; Pages 279-289

<https://doi.org/10.1161/CIRCULATIONAHA.123.066485>




ORIGINAL RESEARCH ARTICLE

Safety of Switching From a Vitamin K Antagonist to a Non-Vitamin K Antagonist Oral Anticoagulant in Frail Older Patients With Atrial Fibrillation: Results of the FRAIL-AF Randomized Controlled Trial

- 8 Dutch thrombosis clinics
- Nonvalvular AFib patients ≥ 75 yo on VKA, eGFR ≥ 30 ml/min
- Groningen frailty score ≥ 3
- Randomized, open label
 - Continue VKA (n=661)
 - Switch to DOAC (n=662)
- Superiority trial with planned interim analysis at 160 events



Cumulative incidence of bleeding DOACs vs VKA



FRAIL-AF: Bleeding vs Thrombosis

Variable	Switch to NOAC		Continue with VKA		Hazard ratio (95% CI)
	n (%)	No. of events/100 patient-years (95% CI)	n (%)	No. of events/100 patient-years (95% CI)	
Primary outcome					
Major or CRNM bleeding	101 (15.3)	17.8 (14.5–21.6)	62 (9.4)	10.5 (8.0–13.4)	1.69 (1.23–2.32)
Secondary outcomes					
Bleeding outcomes separately					
Major bleeding	24 (3.6)	3.9 (2.5–5.9)	16 (2.4)	2.6 (1.5–4.2)	1.52 (0.81–2.87)
CRNM bleeding	84 (12.7)	14.6 (11.7–18.1)	49 (7.4)	8.2 (6.1–10.9)	1.77 (1.24–2.52)
Thromboembolic events	16 (2.4)	2.6 (1.5–4.3)	13 (2.0)	2.1 (1.1–3.6)	1.26 (0.60–2.61)
Composite of thromboembolic events plus major or CRNM bleeding	115 (17.4)	20.6 (17.0–24.7)	73 (11.0)	12.4 (9.8–15.6)	1.65 (1.23–2.21)
Composite of ischemic and hemorrhagic stroke	14 (2.1)	2.3 (1.3–3.8)	11 (1.7)	1.8 (0.9–3.2)	1.30 (0.59–2.87)
All-cause mortality	44 (6.7)	7.1 (5.2–9.5)	46 (7.0)	7.4 (5.4–9.8)	0.96 (0.64–1.45)



FRAIL-AF: Discussion Points

Table 3. First Major or Clinically Relevant Nonmajor Bleeding* Location per Treatment Arm

Bleeding location	Major bleedings		CRNM bleedings	
	Switch to NOAC	Continue with VKA	Switch to NOAC	Continue with VKA
Skin, n (%)			23 (3.5)	10 (1.5)
Oropharyngeal, n (%)		1 (0.2)	19 (2.9)	16 (2.3)
Gastrointestinal, n (%)	9 (1.4)	1 (0.2)	8 (1.2)	3 (0.5)
Urogenital, n (%)			20 (3.0)	11 (1.7)
Brain,† n (%)	7 (1.1)	6 (0.9)		
Ophthalmic, n (%)		1 (0.2)	3 (0.5)	2 (0.3)
Musculoskeletal, n (%)	1 (0.2)		1 (0.2)	4 (0.6)
Lung, n (%)		1 (0.2)		
Other, n (%)	2 (0.3)	3 (0.5)	8 (1.2)	3 (0.5)

Stopped early for futility
(underpowered)

Precludes drawing conclusions on
differences between the groups

VKA patients already tolerant

DOAC choice not individualized or
randomized (50% rivaroxaban)

TTR not reported for VKA arm
(likely >65-70% given setting)



Comorbidities to consider when choosing anticoagulant

- Renal insufficiency
 - dose reductions or avoidance for some DOACs
- Liver disease
 - Caution with VKA and DOACs based on Child-Pugh
- Underweight
 - Dose reduction with Apixaban/Edoxaban in AFib
- Cancer-associated VTE
 - Apixaban/Rivaroxaban/Edoxaban/LMWH > VKA
- Antiphospholipid syndrome
 - VKA > DOACs
- Mechanical Heart valves
 - VKA > DOACs

Comorbidities to consider when choosing anticoagulant


- Altered GI anatomy or enteral nutrition
 - Unpredictable absorption of OAC

DOACs for Enteral Administration

DOAC	Bioavailability	Safe to Crush?	Administer per NG/G Tube?
Apixaban	~50% (unaffected by food)	Yes	Yes
Rivaroxaban	>90% with food† (~66% without food)	Yes	Yes*
Dabigatran	~5% (unaffected by food)	No	No


†Doses ≤10 mg can be administered without regard to food

*Avoid administering post-pyloric



J. Chohan

Geriatric Syndromes: Intersectionality of Multiple Conditions/Issues



Most older AF patients have ≥ 1 geriatric syndrome

\uparrow # geriatric syndromes associated with \downarrow anticoagulant use

May be a key driver of undertreatment of AF

Should be incorporated into decisions around stopping AC



Key Considerations in Choosing an Anticoagulant WITH Your Older Patient

Switching from VKA to DOAC might be associated with similar benefits regarding major bleeds, stroke and mortality.

However, it may be associated with a higher rate of GI bleeding.



Weigh potential gains of simpler treatment with potential increased risk of GI bleeding.

Most important is maintaining long-term adherence, whether on VKA or DOAC.



Case 4: Audience Response Question

- **96 yo** woman on apixaban for atrial fibrillation
- Brought into ED from long-term care facility after a **fall event** that she does not remember
- Has **advanced Alzheimer's** and is **fully dependent** for ADLs
- Patient **intermittently refuses** oral medications at long-term facility
- Head CT is negative for any bleeding and ED resident is asking for recommendations on resuming apixaban

What would you recommend?

- A. Continue twice daily apixaban
- B. Switch to once-daily rivaroxaban
- C. Switch to VKA
- D. Stop all anticoagulation



Stopping Anticoagulants: Need for a Patient-Centered Framework

Competing risk of death from non-stroke causes, such as advanced dementia, diminishes the net clinical benefit (NCB) of anticoagulant therapy

After age 87 years and 92 years, NCB of warfarin and apixaban, respectively, falls below the minimal clinically relevant threshold

Recent data suggests roughly 1/3 of nursing home residents with AF and advanced dementia remain on anticoagulation in last 6 months of life

More high-quality data is needed to inform decision-making and drive antithrombotic stewardship initiatives in these patient populations

“Drive to Deprescribe” initiative (<https://paltc.org/drive-deprescribe>)


Ouellet GM, et al. JAMA Intern Med. 2021;181(8):1121-1123.

Shah SJ, et al. Circ Cardiovasc Qual Outcomes. 2019 Nov;12(11):e006212

Parks A, et al. JAMA Intern Med. 2021;181(8):1123.




Key Considerations in Stopping an Anticoagulant WITH Your Older Patient




Antithrombotic Stewardship: Drive to Deprescribe (D2D) Initiative

- Post-Acute and Long-term Care Med Association (PALTmed) and American Society of Consultant Pharmacists (ASCP)
- Reduce polypharmacy and enhancing patient care
- Deprescribe ASA



Risk Factors ↑ Bleeding:
-Age
-Male
-Diabetes
-Smoking
-↑ Blood pressure
-NSAID use
-History of GI ulcer

Risk Factors ↓ Bleeding:
-PPI use



Practical considerations when managing anticoagulation in older patients

Checklist of clinical parameters to assess when initiating AC and during follow-up

- ✓ Assess Thrombotic Risk (CHA₂DS₂-Vasc or VTE recurrence)
- ✓ Assess bleeding risk using RAMs and other risks specific to elderly
- ✓ Check baseline labs- renal function, coagulation tests, liver function, CBC
- ✓ Check for CI to DOACs or VKA. Alternatives to AC e.g. LAAO
- ✓ Choose suitable AC and dose (QD vs BID) and dose reductions
- ✓ Identify potential drug-drug interactions
- ✓ Reduce unnecessary medications (ASA and others with antiplatelet properties)
- ✓ Evaluate need for PPI
- ✓ Tailor patient education to elderly and repeat as needed. Consider additional teaching/adherence tools
- ✓ Provide anticoagulation card/medication alert tag
- ✓ Determine appropriate frequency of follow-up (early initial F/U may be required in older patients)
- ✓ Explore reasons for nonadherence
- ✓ Explore options for dose reduction or discontinuation of anticoagulation



Anticoagulant Alert Card

This patient is taking anticoagulant therapy.
This card should be carried at all times and shown to healthcare professionals.

Name of patient:	
Address:	
Postcode:	Telephone:
Name of next of kin:	
Hospital number:	NHS Number:

Details of anticoagulant therapy:

Name of anticoagulant:	
Indication for treatment:	
Therapeutic range (INR):	
Treatment started:	Duration of treatment:
Name and address of anticoagulant clinic:	
Telephone number of clinic:	

Acknowledgements

Thanks to the Anticoagulation Forum for slides from a recent webinar on Anticoagulation in Older Patients

<https://acforum.org/web/education-webinars.php>